

#### **4-Quadranten-Rehabilitation nach parodontal-funktioneller und kariöser Schädigung**

**Dr. Sven Egger, MSc, MSc, Prof. Dr.med.dent. Markus Greven, MSc, MDS, PhD**

**DI (27) 1/2023, 13 – 19**

- [1] Joran R., Bodechtel C., Hertrampf K., Hoffmann T., Kocher T., Nitschke I., Schiffner U., Stark H., Zimmer S., Micheelis W., DSM V Surveillance Investigators' Group The Fifth German Oral Health Study (Fünfte Deutsche Mundgesundheitsstudie, DMS V) – rationale, design, and methods. *BMC Oral Health* 2014; 14: 161., <https://pubmed.ncbi.nlm.nih.gov/25547464>.
- [2] Witkowski S, Schicha K. Prep Veneers & Non-Prep Veneers. Berlin, Quintessenz, 2010.
- [3] Egger S, Greven M. Funktionell-implantologische Vier-Quadranten-Rehabilitation (Fallstudie). *Praktische Implantologie und Implantatprothetik (PIP)* 2021; 12 (1): 8-15.
- [4] Egger S, Berg C. Four-quadrant rehabilitation after periodontal functional and erosive damage. *Journal of craniomandibular function* 2019; 11 (1): 75-90.
- [5] D'Amico A. The canine teeth-normal functional relation of the natural teeth of man. *J South Calif Dent Assoc.* 1958; 26: 6-23, 49-60, 127-142, 175-182, 194-208, 239-241.
- [6] Keshvad A, Winstanley RB. An appraisal of the literature on centric relation. Part I. *Int J Oral Rehabil.* 2000a; 27: 823-833.
- [7] Ash MM, Nelson SJ. Wheelers Dental Anatomy, Physiology and Occlusion. 8th ed. Saunders, Philadelphia, 2003 : 417, 419.
- [8] Greven M, Otsuka T, Zutz L, Weber B, Elger C, Sato S. The amount of TMJ displacement correlates with brain activity. *Cranio.* 2011; 29 (4): 291–6.
- [9] Otsuka T, Sato S, et al. Effects of mandibular deviation on brain activation during clenching: an fMRI study. *J Cran Mand Prac.* 2009; 27 (2).
- [10] Levy JH. Teeth as Sensory Organs. *VISTAS – Dawson.* 2009; 2 (3).
- [11] Egger S, Greven M. Funktionell-implantologische Vier-Quadranten-Rehabilitation (Fallstudie). *Praktische Implantologie und Implantatprothetik (PIP)* 2021; 12 (1):8-15.
- [12] Magne P, Belser U. Adhäsig befestigte Keramikrestaurationen. Berlin, Quintessenz, 2004.
- [13] Rufenacht CR. Fundamentals of Esthetics. Chicago, Quintessenz, 1990: 67-134.
- [14] Zuchelli G, De Sanchis M. Cervical abrasion associated with gingival recession: a treatment-based classification system. *J Periodontol.* 2014; (in press).
- [15] Lindhe J, Nyman S. Long-term maintenance of patients treated for advanced periodontal disease. *J Clin Periodontol.* 1984; 11: 504.
- [16] Abduo J. *Quintessence Int.* 2012; 43 (5): 369-80. [www.ncbi.nlm.nih.gov/pubmed/22536588](http://www.ncbi.nlm.nih.gov/pubmed/22536588).
- [17] Gutowski A. Die systematische Behandlung des Abrasionsgebisses von Prof. AlexanderGutowski. Lehrfilm: Live-Demonstrationskurs mit Patienten, Alexander Gutowski, 2010.
- [18] Vailati F, Belser UC. Full-mouth adhesive rehabilitation of a severely eroded dentition: the three-step technique. Part 1. *Eur J Esthet Dent.* 2008; 3 (1): 30-44.
- [19] Vailati F, Belser UC. Full-mouth adhesive rehabilitation of a severely eroded dentition: the three-step technique. Part 2. *Eur J Esthet Dent.* 2008; 3 (2): 128-4.
- [20] Vailati F, Belser UC. Full-mouth adhesive rehabilitation of a severely eroded dentition: the three-step technique. Part 3. *Eur J Esthet Dent.* 2008; 3 (3): 236-57.
- [21] Tarnow DP, Cho SC, Wallace SS. The effect of interimplant distance on the height of inter-implant bone crest. *J Periodontol.* 2000; 71: 546-549.
- [22] Greven M, Seuser W, Pogodin D, Egger S. Full-Mouth-Rehabilitation im funktionsgestörten Erosions- und Abrasionsgebiss, Teil 1: Einfluss der statischen und dynamischen Okklusion auf das Kausystem. *Cosmetic Dentistry* 2020; 18 (2+3): 16-21.

[23] Greven M, Seuser W, Pogodin D, Egger S. Full-Mouth-Rehabilitation im funktionsgestörten Erosions- und Abrasionsgebiss, Teil 2: Bedeutung der Okklusion mit Fallbeispiel. Cosmetic Dentistry 2020; 18 (4): 16-21.