

## Literatur

# Next Level – moderne Tissue-Level-Implantate

PD Dr. med. Dr. med. dent. Eik Schiegnitz, M.Sc.

### DENTALE IMPLANTOLOGIE & PARODONTOLOGIE, Juni 2021

- [1] Cochran D, Stavropoulos A, Obrecht M, Pippenger B, Dard M A Comparison of Tapered and Nontapered Implants in the Minipig. *Int J Oral Maxillofac Implants* 2016, 31(6):1341-1347. doi: 10.11607/jomi.4712.
- [2] Derks J, Håkansson J, Therapy Wennström JL, Tomasi C, Larsson M, Berglundh T. Effectiveness of Implant Analyzed in a Swedish Population: Early and late Implant Loss. *J Dent Res* 2014, Dec 11.
- [3] Derks J, Schaller D, Håkansson J, Wennström JL, Tomasi C, Berglundh T. Effectiveness of Implant Therapy Analyzed in a Swedish Population: Prevalence of Peri-implantitis. *J Dent Res*. 2016, 95(1):43-9. doi: 10.1177/0022034515608832. PMID: 26701919.
- [4] Gottlow J, Dard M, Kjellson F, Obrecht M, Sennerby L. Evaluation of a new titanium-zirconium dental implant: a biomechanical and histological comparative study in the mini pig. *Clin Implant Dent Relat Res*. 2012 Aug;14(4):538-45.
- [5] Kang MH, Jung UW, Cho KS, Lee JS. Retrospective radiographic observational study of 1692 Straumann tissue-level dental implants over 10 years. II. Marginal bone stability. *Clin Implant Dent Relat Res* 2018, Epub Jul 26, doi: 10.1111/cid.12636.
- [6] Kim S, Jung UW, Cho KS, Lee JS. Retrospective radiographic observational study of 1692 Straumann tissue-level dental implants over 10 years: I. Implant survival and loss pattern. *Clin Implant Dent Relat Res* 2018, Epub Aug 15, doi: 10.1111/cid.12659.
- [7] Kopf BS, Ruch S, Berner S, Spencer ND, Maniura-Weber K. The role of nanostructures and hydrophilicity in osseointegration: In-vitro protein-adsorption and blood-interaction studies. *J Biomed Mater Res A* 2015, 103(8):2661-72. doi: 10.1002/jbm.a.35401. Epub 2015, Feb 11.
- [8] Lang NP, Salvi GE, Huynh-Ba G, Ivanovski S, Donos N, Bosshardt DD. Early osseointegration to hydrophilic and hydrophobic implant surfaces in humans. *Clin Oral Implants Res* 2011, 22(4):349-56. doi: 10.1111/j.1600-0501.2011.02172.x.
- [9] Pecanov-Schröder A: Schmale konische Implantate: Schnelle Osseointegration und hohe prothetische Flexibilität. *Die ZahnarztWoche DZW OI* 2017;17 (Beilage Orale Implantologie 1):36-38.

- [10] Rupp F, Scheideler L, Olshanska N, de Wild M, Wieland M, Geis-Gerstorfer J. Enhancing surface free energy and hydrophilicity through chemical modification of microstructured titanium implant surfaces. *J Biomed Mater Res A* 2006, 76(2):323-34.
- [11] Schiegnitz E, Kämmerer PW, Hellwich P, König J, Sagheb K, Al-Nawas B. Treatment concepts of horizontally deficient alveolar ridges - a retrospective clinical study comparing narrow diameter implants (3.3 mm) in pristine bone versus standard diameter implants (4.1 mm) in augmented bone. 2021 (in review)
- [12] Straumann-Gruppe: XenoGraft und XenoFlex: Für Routinefälle, Broschüre. Institut Straumann AG (2019)
- [13] Quirynen M, Al-Nawas B, Meijer HJ, Razavi A, Reichert TE, Schimmel M, Storelli S, Romeo E; Roxolid Study Group: Small-diameter titanium Grade IV and titanium-zirconium implants in edentulous mandibles: three-year results from a double-blind, randomized controlled trial. *Clin Oral Implants Res* 2015, 26(7):831-40. doi: 10.1111/clr.12367. Epub 2014 Apr 9.
- [14] Wen B, Zhu F, Li Z, Zhang P, Dard M: The osseointegration behavior of titanium-zirconium implants in ovariectomized rabbits. *Clin Oral Implants Res* 2014, 25(7):819-25. doi: 10.1111/clr.12141. Epub 2013 Feb 21.