

Literatur

## Eine Alternative zum autogenen Knochen und zur lokalen Lappenplastik?

Prof. Shahram Ghanaati et al.

DENTALE IMPLANTOLOGIE Mai 2018

- [1] Arx T, Kurt B. Implant placement and simultaneous ridge augmentation using autogenous bone and a micro titanium mesh: a prospective clinical study with 20 implants. *Clin Oral Implants Res.* 1999;10(1):24-33.
- [2] Atwood DA. Reduction of residual ridges: a major oral disease entity. *J Prosthet Dent.* 1971;26(3):266-279.
- [3] Barbeck M, Lorenz J, Kubesch A, et al. Porcine Dermis-Derived Collagen Membranes Induce Implantation Bed Vascularization Via Multinucleated Giant Cells: A Physiological Reaction? *J Oral Implantol.* 2015;41(6):e238-51.
- [4] Barbeck M, Udeabor SE, Lorenz J, et al. Induction of multinucleated giant cells in response to small sized bovine bone substitute (Bio-Oss<sup>TM</sup>) results in an enhanced early implantation bed vascularization. *Ann Maxillofac Surg.* 2014;4(2):150-157.
- [5] Boudet C. Platelet-rich fibrin in mesh exposure repair. *Dent Today.* 2014;33(1):112-113.
- [6] Cawood JI, Howell RA. A classification of the edentulous jaws. *Int J Oral Maxillofac Surg.* 1988;17(4):232-236.
- [7] Choukroun J, Ghanaati S. Reduction of relative centrifugation force within injectable platelet-rich-fibrin (PRF) concentrates advances patients' own inflammatory cells, platelets and growth factors: the first introduction to the low speed centrifugation concept. *Eur J Trauma Emerg Surg.* 2018;44(1):87-95.
- [8] Cucchi A, Vignudelli E, Napolitano A, Marchetti C, Corinaldesi G. Evaluation of complication rates and vertical bone gain after guided bone regeneration with non-resorbable membranes versus titanium meshes and resorbable membranes. A randomized clinical trial. *Clin Implant Dent Relat Res.* 2017;19(5):821-832.
- [9] DiGiovanni CW, Petricek JM. The evolution of rhPDGF-BB in musculoskeletal repair and its role in foot and ankle fusion surgery. *Foot Ankle Clin.* 2010;15(4):621-640.
- [10] Elgali I, Turri A, Xia W, et al. Guided bone regeneration using resorbable membrane and different bone substitutes: Early histological and molecular events. *Acta Biomater.* 2016;29:409-423.
- [11] Forsberg S, Rollman O. Re-epithelialization from human skin explant cultures is promoted by ligand-activated HER3 receptor. *J Dermatol Sci.* 2010;59(1):7-15.
- [12] Ghanaati S, Barbeck M, Detsch R, et al. The chemical composition of synthetic bone substitutes influences tissue reactions in vivo : histological and histomorphometrical analysis of the cellular inflammatory response to hydroxyapatite, beta-tricalcium phosphate and biphasic calcium phosphate ce. *Biomed Mater.* 2012;7(1):15005.
- [13] Ghanaati S, Barbeck M, Lorenz J, et al. Synthetic bone substitute material comparable with xenogeneic material for bone tissue regeneration in oral cancer patients: First and preliminary histological, histomorphometrical and clinical results. *Ann Maxillofac Surg.* 2013;3(2):126-138.

- [14]Ghanaati S, Booms P, Orlowska A, et al. Advanced Platelet-Rich Fibrin: A New Concept for Cell-Based Tissue Engineering by Means of Inflammatory Cells. *J Oral Implantol.* 2014;40(6):679-689.
- [15]Ghanaati S, Schlee M, Webber MJ, et al. Evaluation of the tissue reaction to a new bilayered collagen matrix in vivo and its translation to the clinic. *Biomed Mater.* 2011;6(1):15010.
- [16]Ghanaati S. Non-cross-linked porcine-based collagen I-III membranes do not require high vascularization rates for their integration within the implantation bed: A paradigm shift. *Acta Biomater.* 2012;8(8):3061-3072.
- [17]Ghanaati S. et. al., Komplexe implantologische Rehabilitiation mittels eines dreidimensionalen Titangitters, autogenem Fibrikkonzentrat (PRF) und eines xenogenem Knochenersatzmaterials, *Dent. Implant.* 20, 6, 368 - 375 (2016),
- [18]Khoury F, D oliveux R. The Bone Core Technique for the Augmentation of Limited Bony Defects: Five-Year Prospective Study with a New Minimally Invasive Technique. *Int J Periodontics Restorative Dent.* 38(2):199-207.
- [19]Krause, S Möglichkeiten und Grenzen komplexer Augmentationen mittels CAD/CAM gefertigter Titangitter, 2017 dentale implatologie und parodontologie
- [20]Lorenz J, Kubesch A, Al-Maawi S, et al. Allogeneic bone block for challenging augmentation—a clinical, histological, and histomorphometrical investigation of tissue reaction and new bone formation. *Clin Oral Investig.* 2018. doi:10.1007/s00784-018-2407-0.
- [21]Lorenz J, Kubesch A, Korzinskas T, et al. TRAP-Positive Multinucleated Giant Cells Are Foreign Body Giant Cells Rather Than Osteoclasts: Results From a Split-Mouth Study in Humans. *J Oral Implantol.* 2015;41(6):e257-e266.
- [22]M. Seiler et al. Augmentation with a CAD/CAM Titanium Mesh in the Anterior Mandible, , *Clinical Science* 1, 2016, S. 2-9
- [23]M. Seiler et al.CAD/CAM Lösung für die Augmentation mit CAD/CAM-gefertigtem Titangitter in der Unterkieferfront, , *J. Cont. Dent. Education*, 2016, 1, 2-9
- [24]Moens S, Goveia J, Stapor PC, Cantelmo AR, Carmeliet P. The multifaceted activity of VEGF in angiogenesis – Implications for therapy responses. *Cytokine Growth Factor Rev.* 2014;25(4):473-482.
- [25]Sagheb K, Schiegnitz E, Moergel M, Walter C, Al-Nawas B, Wagner W. Clinical outcome of alveolar ridge augmentation with individualized CAD-CAM-produced titanium mesh.
- [26]Sakkas A, Wilde F, Heufelder M, Winter K, Schramm A. Autogenous bone grafts in oral implantology-is it still a "gold standard"? A consecutive review of 279 patients with 456 clinical procedures. *Int J Implant Dent.* 2017;3(1):23. doi:10.1186/s40729-017-0084-4.
- [27]Schneider D, Weber FE, Grunder U, Andreoni C, Burkhardt R, Jung RE. A randomized controlled clinical multicenter trial comparing the clinical and histological performance of a new, modified polylactide-co-glycolide acid membrane to an expanded polytetrafluoroethylene membrane in guided bone regeneration procedures. *Clin Oral Implants Res.* 2014;25(2):150-158.
- [28]Sumida T, Otawa N, Kamata YU, et al. Custom-made titanium devices as membranes for bone augmentation in implant treatment: Clinical application and the comparison with conventional titanium mesh. *J Craniomaxillofac Surg.* 2015;43(10):2183-2188.
- [29]Weibull L, Widmark G, Ivanoff C-J, Borg E, Rasmusson L. Morbidity after Chin Bone Harvesting - A Retrospective Long-Term Follow-Up Study. *Clin Implant Dent Relat Res.* 2009;11(2):149-157.
- [30]Wend S, Kubesch A, Orlowska A, et al. Reduction of the relative centrifugal force influences cell number and growth factor release within injectable PRF-based matrices. *J Mater Sci Mater Med.*