

Literatur

Das periimplantäre Weichgewebe als Schlüsselfaktor für einen Langzeiterfolg

Dr. Jan Klenke

DENT IMPLANTOL 21, 6, 380-383 (2017)

- [1] Mehdi A, et al.: Significance of keratinized mucosa on the health status of the supporting tissue around implants supporting overdentures. *Journal of oral Implantology*. 2009. 35 (5): 323 – 237
- [2] Berglundh T., et al.: The soft tissue barrier at implants and teeth. *Clinical oral Implants research*. 1991. 2: 81 – 90.
- [3] Berglundh T., et al.: Soft tissue reactions to de novo plaque formation at implants and teeth. An experimental study in the dog. *Clinical oral Implants Research*. 1992. 3: 1 – 8.
- [4] Berglundh T., et al.: Dimension of the peri-implant mucosa. Biological width revisited. *Journal of Clinical Periodontology*. 1996. 23: 971 – 973.
- [5] Ericsson I., et.al.: Long standing plaque and gingivitis at implants and teeth in the dog. *Clinical Oral Implants Research*. 1992. 3: 99 – 103.
- [6] Giovanni B., et al.: Clinical study of a flap design to increase the keratinized gingiva around implants: 4 year follow up. *Journal of Oral Implantology*. 2014. 40 (4): 459 – 464
- [7] Wennstrom JL, Derks J: Is there a need for keratinized mucosa around implants to maintain health and tissue stability? *Clin Oral Implants Res* 2012; 23 (6 Suppl): 136-146.)
- [8] Adell R, Lekholm U, Rockler B, Branemark PI, Lindhe J, Eriksson B, Sbordone L.: Marginal reactions at osseointegrated titanium fixtures. A 3-year longitudinal prospective study. *Int J Oral Maxillofac surg* 1986; 15: 39-52.
- [9] Adibrad M, Shahabuei M, Sahabi M: Significance of the width of keratinized mucosa on the health status of the supporting tissue around implants supporting overdentures. *J Oral Implantol* 2009; 35: 232-237.)
- [10] Linkevicius T, Apse P, Grybauskas S, Puisys A: Reaction of crestal bone around implants depending on mucosal tissue thickness. A 1-year prospective clinical study. *Stomatologica* 2009;11: 83-91.