

Literatur

Implantate mit hoher Primärstabilität erleichtern Knochengewinn

Fabian Hirsch

- [1] Altıparmak N, Akdeniz SS, Bayram B, Gulsever S, Uçkan S: Alveolar ridge splitting versus autogenous onlay bone grafting: Complications and implant survival rates. *Implant Dent* 26(2): 284-287 (2017).
- [2] Bassetti MA, Bassetti RG, Bosshardt DD: The alveolar ridge splitting/expansion technique: a systematic review. *Clin Oral Implant Res* 27:310-324 (2016).
- [3] Becktor JP, Starch-Jensen T: Maxillary alveolar ridge expansion with split-crest technique compared with lateral ridge augmentation with autogenous bone block graft: A systematic review. *J Oral Maxillofac Res* 10(4):e2, online am 30. Dezember, doi: 10.5037/jomr.2019.10402 (2019).
- [4] Berger S, Hakl P, Sutter W, Meier M, Roland H, Bandura P, Turhani D: Interantral alveolar ridge splitting for maxillary horizontal expansion and simultaneous dental implant insertion: A case report. *Annals of Medicine and Surgery* 48:83-87 (2019).
- [5] Dohiem MM, Nassar HI, El Charkawi H: Bone changes in ridge split with immediate implant placement: A systematic review. *Future Dental J* 6-12 (2015).
- [6] Le Guéhennec L, Soueidan A, Layrolle P, Amouriq Y: Surface treatments of titanium dental implants for rapid osseointegration. *Dent Mater* 23(7):844-54 (2007).
- [7] Olate S, Marin A, Oporto G, Farias D, Cantin M: Alveolar ridge splitting for implant installation of atrophic sites. Analysis of a case series. *Int J Odontostomat* 9(2):249-254 (2015).