

# Rekonstruktive Behandlung bei Aplasie der lateralen Schneidezähne im Oberkiefer

Dr. Mathias Plöger

- [1] Terhyden H: Implantatversorgung bei Kindern, Jugendlichen und jungen Erwachsenen. *Implantologie* 2018; 26(2):115-122
- [2] Chaushu G, Nissan J: The use of cancellous bone block allograft for reconstruction of the atrophic alveolar ridge in: Haim T (ed): *Bone regeneration*, In-Tech, Rijeka, Croatia 2012
- [3] Björk A, Skieller V: Growth of the maxilla in three dimensions as revealed radiographically by the implant method. *Br J Orthod* 1977; 4: 53-64
- [4] Bernhard JP, Schatz JP, Christou P, Belser U, Kiliardis S. Long-term vertical changes of the anterior maxillary teeth adjacent to single implants in young and mature adults. A retrospective study. *J Clin Periodontol* 2004; 31: 1024-1028
- [5] Buser D, Belser U, Wismeijer D. *Ästhetische Implantattherapie – Einzelzahnersatz*. ITI, Treatment Guide Series Bd. 1. Berlin: Quintessenz, 2007
- [6] Khoury F: *Augmentative Verfahren in der Implantologie*. Berlin: Quintessenz, 2009
- [7] Chiapasco M, Casentini P, Zaniboni M: Bone augmentation procedures in implant dentistry. *Int J Oral Maxillofac Implants* 2009 (suppl): 237-259
- [8] Jensen SS, Terheyden H: Bone augmentation procedures in localized defects in the alveolar ridge: clinical results with different bone grafts and bone-substitute materials. *Int J Oral Maxillofac Implants* 2009; 24 (suppl): 218-236
- [9] Plöger M, Schau I: *Allogene Knochenblöcke in der zahnärztlichen Implantologie*. Spitta Verlag, Balingen 2010
- [10] Aghaloo TL, Moy PK: Which hard tissue augmentation techniques are the most successful in furnishing bony support for implant placement. *Int J oral Maxillofac Implants* 2007; 22 (suppl): 49-70
- [11] Al Nawas B, Schiegnitz E: Augmentation procedures using bone substitute materials or autogenous bone. *EUR J Oral Implantol* 2014; 7 (suppl): 1-16
- [12] Berggren A., Weiland AJ, Dorfman H: Free vascular bone grafts. Factors affecting their survival and ability to heal the recipient bone defects. *Plast Reconstr Surg* 1982; 69: 19-29
- [13] Calori GM, Mazza E, Colombo M, Ripamonti C: The use of bone-graft substitutes in large bone defects: any specific needs? *Injury* 2011; 42 Suppl 2: S56-63
- [14] Dimitriou R, Metaliotakis GI, Angoules AG, Kanakoris MK, Giannoudis PV: Complications following autologous bone graft harvesting from iliac crest and using RIA a systematic review. *Injury* 2011; 43 (suppl 2): 3-15
- [15] Esposito M, Grusovin MG, Felice P, Karatzopoulos G, Worthington HV, Coulthard P: The efficacy of horizontal and vertical bone augmentation procedures for dental implants – a Cochrane systematic review. *Eur J Oral Implantol* 2009; 2: 167-184
- [16] Esser E, Hümmeke S, Krebs M, Maier F: The use of allogeneic bone grafts for pre-implant alveolar ridge augmentation. *Z Zahnärztl Impl* 2016; 32 (4): 284-296
- [17] Nevins ML, Reynolds MA: Tissue engineering with recombinant human platelet-derived growth factor BB for implant site development. *Compend Contin Educ Dent* 2011; 32(2): 30-36
- [18] Nkenke E, Weisbach V, Winckler E, Kessler P, Schultze-Mosgau S, Wilffang J, Neukam FW: Morbidity of harvesting of bone grafts from the iliac crest for preprosthetic augmentation procedures: a prospective study. *Int J Oral Maxillofac Surg* 2004; 33: 157-163
- [19] Pommer B, Zechner W, Watzek G, Palmer R: To graft or not to graft? Evidence-based guide to decision making in oral bone graft surgery. In: Zorzi A (Ed) *Bone grafting*, Intech Europe 2012, Rijeka, Croatia
- [20] Scheerlinck LME, Muradin MSM, Van der Bilt A, Meijer GJ, Koole R, van Cann E: Donor site complication in bone grafting-comparison of iliac, calvarial and mandibular ramus. *Int J Maxillofac Implants* 2013; 28: 221-227
- [21] Schlee M, Rothamel D: Ridge augmentation using customized bone blocks: proof of concept and histologic finding. *Impl Dent* 2013; 22: 212-218
- [22] Tetsch J, Tetsch P, Lysek D: Long-term results after lateral and osteotome technique sinus floor elevation: a retrospective analyses of 2190 implants over a time period of 15 years. *Clin Oral Implants Res* 2010; 21: 497-503